

# CONCUSSION FACT SHEET FOR PARENTS



## WHAT IS A CONCUSSION?

A concussion is a type of traumatic brain injury. Concussions are caused by a bump or blow to the head. Even a "ding," "getting your bell rung," or what seems to be a mild bump or blow to the head can be serious.

You can't see a concussion. Signs and symptoms of concussion can show up right after the injury or may not appear or be noticed until days or weeks after the injury. If your child reports any symptoms of concussion, or if you notice the symptoms yourself, seek medical attention right away.

## WHAT ARE THE SIGNS AND SYMPTOMS OF CONCUSSION?

If your child has experienced a bump or blow to the head during a game or practice, look for any of the following signs of a concussion:

### SYMPTOMS REPORTED BY ATHLETE:

- Headache or "pressure" in head
- Nausea or vomiting
- Balance problems or dizziness
- Double or blurry vision
- Sensitivity to light
- Sensitivity to noise
- Feeling sluggish, hazy, foggy, or groggy
- Concentration or memory problems
- Confusion
- Just not "feeling right" or is "feeling down"

### SIGNS OBSERVED BY PARENTS/ GUARDIANS:

- Appears dazed or stunned
- Is confused about assignment or position
- Forgets an instruction
- Is unsure of game, score, or opponent
- Moves clumsily
- Answers questions slowly
- Loses consciousness (even briefly)
- Shows mood, behavior, or personality changes



[ INSERT YOUR LOGO ]



## DANGER SIGNS

Be alert for symptoms that worsen over time. Your child or teen should be seen in an emergency department right away if s/he has:

- One pupil (the black part in the middle of the eye) larger than the other
- Drowsiness or cannot be awakened
- A headache that gets worse and does not go away
- Weakness, numbness, or decreased coordination
- Repeated vomiting or nausea
- Slurred speech
- Convulsions or seizures
- Difficulty recognizing people or places
- Increasing confusion, restlessness, or agitation
- Unusual behavior
- Loss of consciousness (even a brief loss of consciousness should be taken seriously)

## WHAT SHOULD YOU DO IF YOU THINK YOUR CHILD HAS A CONCUSSION?

1. **SEEK MEDICAL ATTENTION RIGHT AWAY**  
A health care professional will be able to decide how serious the concussion is and when it is safe for your child to return to regular activities, including sports.
2. **KEEP YOUR CHILD OUT OF PLAY.**  
Concussions take time to heal. Don't let your child return to play the day of the injury and until a health care professional says it's OK. Children who return to play too soon - while the brain is still healing - risk a greater chance of having a second concussion. Repeat or later concussions can be very serious. They can cause permanent brain damage, affecting your child for a lifetime.
3. **TELL YOUR CHILD'S COACH ABOUT ANY PREVIOUS CONCUSSION.**  
Coaches should know if your child had a previous concussion. Your child's coach may not know about a concussion your child received in another sport or activity unless you tell the coach.

## HOW CAN YOU HELP YOUR CHILD PREVENT A CONCUSSION OR OTHER SERIOUS BRAIN INJURY?

- Ensure that they follow their coach's rules for safety and the rules of the sport.
- Encourage them to practice good sportsmanship at all times.
- Make sure they wear the right protective equipment for their activity. Protective equipment should fit properly and be well maintained.
- Wearing a helmet is a must to reduce the risk of a serious brain injury or skull fracture.
  - However, helmets are not designed to prevent concussions. There is no "concussion-proof" helmet. So, even with a helmet, it is important for kids and teens to avoid hits to the head.

## HOW CAN I HELP MY CHILD RETURN TO SCHOOL SAFELY AFTER A CONCUSSION?

Children and teens who return to school after a concussion may need to:

- Take rest breaks as needed
- Spend fewer hours at school
- Be given more time to take tests or complete assignments
- Receive help with schoolwork
- Reduce time spent reading, writing, or on the computer

Talk with your child's teachers, school nurse, coach, speech-language pathologist, or counselor about your child's concussion and symptoms. As your child's symptoms decrease, the extra help or support can be removed gradually.



JOIN THE CONVERSATION  [www.facebook.com/CDCHeadsUp](http://www.facebook.com/CDCHeadsUp)

TO LEARN MORE GO TO >> [WWW.CDC.GOV/CONCUSSION](http://WWW.CDC.GOV/CONCUSSION)

Content Source: CDC's Heads Up Program. Created through a grant to the CDC Foundation from the National Operating Committee on Standards for Athletic Equipment (NOCSAE).



# CONCUSSION IN YOUTH SPORTS



## A Fact Sheet for PARENTS

### WHAT IS A CONCUSSION?

A concussion is a type of traumatic brain injury. Concussions are caused by a bump or blow to the head. Even a “ding,” “getting your bell rung,” or what seems to be a mild bump or blow to the head can be serious.

You can’t see a concussion. Signs and symptoms of concussion can show up right after the injury or may not appear or be noticed until days or weeks after the injury. If your child reports any symptoms of concussion, or if you notice the symptoms yourself, seek medical attention right away.

### WHAT ARE THE SIGNS AND SYMPTOMS OF A CONCUSSION?

#### Signs Observed by Parents or Guardians

*If your child has experienced a bump or blow to the head during a game or practice, look for any of the following signs and symptoms of a concussion:*

- Appears dazed or stunned
- Is confused about assignment or position
- Forgets an instruction
- Is unsure of game, score, or opponent
- Moves clumsily
- Answers questions slowly
- Loses consciousness (even briefly)
- Shows mood, behavior, or personality changes

#### Symptoms Reported by Athlete

- Headache or “pressure” in head
- Nausea or vomiting
- Balance problems or dizziness
- Double or blurry vision
- Sensitivity to light
- Sensitivity to noise
- Feeling sluggish, hazy, foggy, or groggy
- Concentration or memory problems
- Confusion
- Just “not feeling right” or “feeling down”

### HOW CAN YOU HELP YOUR CHILD PREVENT A CONCUSSION OR OTHER SERIOUS BRAIN INJURY?

- Ensure that they follow their coach’s rules for safety and the rules of the sport.
- Encourage them to practice good sportsmanship at all times.
- Make sure they wear the right protective equipment for their activity. Protective equipment should fit properly and be well maintained.
- Wearing a helmet is a must to reduce the risk of a serious brain injury or skull fracture.
  - However, helmets are not designed to prevent concussions. There is no “concussion-proof” helmet. So, even with a helmet, it is important for kids and teens to avoid hits to the head.

### WHAT SHOULD YOU DO IF YOU THINK YOUR CHILD HAS A CONCUSSION?

#### 1. SEEK MEDICAL ATTENTION RIGHT AWAY.

A health care professional will be able to decide how serious the concussion is and when it is safe for your child to return to regular activities, including sports.

#### 2. KEEP YOUR CHILD OUT OF PLAY.

Concussions take time to heal. Don’t let your child return to play the day of the injury and until a health care professional says it’s OK. Children who return to play too soon—while the brain is still healing—risk a greater chance of having a repeat concussion. Repeat or later concussions can be very serious. They can cause permanent brain damage, affecting your child for a lifetime.

#### 3. TELL YOUR CHILD’S COACH ABOUT ANY PREVIOUS CONCUSSION.

Coaches should know if your child had a previous concussion. Your child’s coach may not know about a concussion your child received in another sport or activity unless you tell the coach.

***It’s better to miss one game than the whole season.***

For more information, visit [www.cdc.gov/Concussion](http://www.cdc.gov/Concussion).



# CONCUSSION IN YOUTH SPORTS

## Hoja Informativa para los PADRES

### ¿QUÉ ES LA CONMOCIÓN CEREBRAL?

Una conmoción cerebral es una lesión en el cerebro, causada por un golpe en la cabeza o una sacudida. Incluso una pequeña conmoción o lo que parece ser un golpe o sacudida leve puede ser serio.

La conmoción cerebral no puede verse. Los signos y síntomas de una conmoción pueden aparecer inmediatamente después de la lesión o puede que no aparezcan, o se hagan visibles algunos días o meses después de haber sufrido la lesión. Si su hijo tiene los signos de una conmoción cerebral o si usted nota algún síntoma, busque atención médica de inmediato.

### ¿CUÁLES SON LOS SIGNOS Y SÍNTOMAS DE LA CONMOCIÓN CEREBRAL?

#### Signos que notan los padres y los tutores

*Si su hijo ha sufrido un golpe en la cabeza o una sacudida durante un juego o una práctica, obsérvelo para determinar si tiene alguno de los siguientes signos y síntomas de una conmoción cerebral:*

- Luce aturdido o fuera de control
- Se confunde con la actividad asignada
- Olvida las jugadas
- No se muestra seguro del juego, la puntuación ni de sus adversarios
- Se mueve con torpeza
- Responde con lentitud
- Pierde el conocimiento (así sea momentáneamente)
- Muestra cambios de conducta o de personalidad
- No puede recordar lo ocurrido antes de un lanzamiento o un caída
- No puede recordar lo ocurrido después de un lanzamiento o un caída

#### Síntomas que reporta el atleta

- Dolor o "presión" en la cabeza
- Náuseas o vómitos
- Problemas de equilibrio, mareo
- Visión doble o borrosa
- Sensibilidad a la luz y al ruido
- Se siente débil, confuso, aturdido o grogui
- Problemas de concentración o memoria
- Confusión
- No se "siente bien"

### ¿CÓMO AYUDAR A SU HIJO A PREVENIR UNA CONMOCIÓN CEREBRAL?

Aunque todo deporte es diferente, hay medidas que puede tomar para protegerse.

- Haga que siga las reglas impartidas por el entrenador y las reglas del deporte que practica.
- Invítelo a mantener el espíritu deportivo en todo momento.
- Haga que su hijo use el equipo protector adecuado según la actividad que realiza. El equipo de protección debe ajustarse bien, debe hacersele el mantenimiento adecuado, y el jugador debe usarlo correctamente y en todo momento.

### ¿QUÉ DEBE HACER SI CREE QUE SU HIJO HA SUFRIDO UNA CONMOCIÓN CEREBRAL?

- 1. Busque atención médica de inmediato.** Un profesional de la salud podrá determinar la seriedad de la conmoción cerebral que ha sufrido el niño y cuándo podrá regresar al juego sin riesgo alguno.
- 2. No permita que su hijo siga jugando.** Las conmociones cerebrales necesitan de un cierto tiempo para curarse. No permita que su hijo regrese al juego hasta que un profesional de la salud le haya dicho que puede hacerlo. Los niños que regresan al juego antes de lo debido—mientras el cerebro está en proceso de curación—corren un mayor riesgo de sufrir otra conmoción. Las conmociones cerebrales siguientes pueden ser muy serias. Pueden causar daño cerebral permanente que afectarán al niño de por vida.
- 3. Informe al entrenador del niño sobre cualquier conmoción cerebral que el niño haya sufrido recientemente.** Los entrenadores deben saber si el niño ha sufrido una conmoción recientemente en CUALQUIER deporte. El entrenador no necesariamente sabrá si el niño ha tenido una conmoción en otro deporte o actividad a menos que usted se lo diga.

***Es preferible perderse un juego que toda la temporada.***

Para obtener más información, visite [www.cdc.gov/ConcussionInYouthSports](http://www.cdc.gov/ConcussionInYouthSports).